

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 875 569 A1

R0058A - IDS
File

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
04.11.1998 Bulletin 1998/45

(51) Int Cl.⁶: C12N 15/12, C07K 14/47

(21) Application number: 98302815.0

(22) Date of filing: 09.04.1998

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(71) Applicant: SMITHKLINE BEECHAM
CORPORATION
Philadelphia Pennsylvania 19103 (US)

(72) Inventor: Feld, John
King of Prussia, Pennsylvania 19406 (US)

(30) Priority: 28.04.1997 US 44974 P
23.09.1997 US 935433

(74) Representative: Crump, Julian Richard John et al
1J Cleveland,
40-43 Chancery Lane
London WC2A 1JQ (GB)

(54) A human sodium dependent phosphate transporter (IPT-1)

(57) IPT-1 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing IPT-1 polypeptides and polynucle-

otides in the design of protocols for the treatment of chronic renal failure, end stage renal disease, uremic bone disease, and cancer, among others, and diagnostic assays for such conditions.

EP 0 875 569 A1

Description

This application claims the benefit of U.S. Provisional Application No: 60/044,974, filed April 28, 1997.

5 FIELD OF INVENTION

This invention relates to newly identified polynucleotides, polypeptides encoded by them and to the use of such polynucleotides and polypeptides, and to their production. More particularly, the polynucleotides and polypeptides of the present invention relate to the sodium dependent phosphate transporters family, hereinafter referred to as IPT-1.
 10 The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

BACKGROUND OF THE INVENTION

Phosphate retention has been shown to play a critical role in the development of uremic bone disease. Blockade of intestinal absorption of phosphate could provide an important target for prevention of uremic bone disease in patients who have end stage renal disease (ESRD) and possibly a target for slowing the progression of renal disease itself. Patients with ESRD cannot excrete phosphate, and they develop hyperphosphatemia, secondary hyperparathyroidism and uremic bone disease. Current treatment of these patients involves dietary phosphate restriction and phosphate binders, both of which have severe drawbacks. Blockade of phosphate absorption with a specific inhibitor of the intestinal phosphate transporter would provide a major advance in the treatment of these patients. This indicates that the sodium dependent phosphate transporters family has an established, proven history as therapeutic targets. Clearly there is a need for identification and characterization of further members of the sodium dependent phosphate transporters family which can play a role in preventing, ameliorating or correcting dysfunctions or diseases, including, but not limited to, chronic renal failure, end stage renal disease, uremic bone disease, and cancer.

SUMMARY OF THE INVENTION

In one aspect, the invention relates to IPT-1 polypeptides and recombinant materials and methods for their production. Another aspect of the invention relates to methods for using such IPT-1 polypeptides and polynucleotides. Such uses include the treatment of chronic renal failure, end stage renal disease, uremic bone disease, and cancer, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with IPT-1 imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate IPT-1 activity or levels.

DESCRIPTION OF THE INVENTION**Definitions**

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"IPT-1" refers, among others, generally to a polypeptide having the amino acid sequence set forth in SEQ ID NO: 2 or an allelic variant thereof.

"IPT-1 activity or IPT-1 polypeptide activity" or "biological activity of the IPT-1 or IPT-1 polypeptide" refers to the metabolic or physiologic function of said IPT-1 including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said IPT-1.

"IPT-1 gene" refers to a polynucleotide having the nucleotide sequence set forth in SEQ ID NO: 1 or allelic variants thereof and/or their complements.

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

"Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide" generally refers to any polyribonucleotide or polydeoxiribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide"

refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Posttranslational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62.

"Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques. See, e.g.: (COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A.M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D.W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinje, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., *et al.*, *Nucleic Acids Research* (1984)

12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. *et al.*, *J Molec Biol* (1990) 215:403).

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% "identity" to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

Polypeptides of the Invention

In one aspect, the present invention relates to IPT-1 polypeptides (or IPT-1 proteins). The IPT-1 polypeptides include the polypeptide of SEQ ID NOS:2 and 4; as well as polypeptides comprising the amino acid sequence of SEQ ID NO: 2; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Also included within IPT-1 polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and still more preferably at least 95% identity to SEQ ID NO:2. Furthermore, those with at least 97-99% are highly preferred. Preferably IPT-1 polypeptide exhibit at least one biological activity of IPT-1.

The IPT-1 polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

Fragments of the IPT-1 polypeptides are also included in the invention. A fragment is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned IPT-1 polypeptides. As with IPT-1 polypeptides, fragments may be "free-standing" or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of IPT-1 polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of IPT-1 polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate IPT-1 activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

Preferably, all of these polypeptide fragments retain the biological activity of the IPT-1, including antigenic activity. Among the most preferred fragment is that having the amino acid sequence of SEQ ID NO: 4. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents

by conservative amino acid substitutions -- i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

The IPT-1 polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

Another aspect of the invention relates to IPT-1 polynucleotides. IPT-1 polynucleotides include isolated polynucleotides which encode the IPT-1 polypeptides and fragments, and polynucleotides closely related thereto. More specifically, IPT-1 polynucleotide of the invention include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding a IPT-1 polypeptide of SEQ ID NO: 2, and polynucleotides having the particular sequences of SEQ ID NOS:1 and 3. IPT-1 polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the IPT-1 polypeptide of SEQ ID NO:2, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO: 1 over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under IPT-1 polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO: 1 to hybridize under conditions useable for amplification or for use as a probe or marker. The invention also provides polynucleotides which are complementary to such IPT-1 polynucleotides.

IPT-1 of the invention is structurally related to other proteins of the sodium dependent phosphate transporters family, as shown by the results of sequencing the cDNA encoding human IPT-1. The cDNA sequence of SEQ ID NO: 1 contains an open reading frame (nucleotide number 64 to 2136) encoding a polypeptide of 691 amino acids of SEQ ID NO:2. The amino acid sequence of Table 1 (SEQ ID NO:2) has about 76 identity (using FASTA) in 691 amino acid residues with bovine sodium dependent phosphate transporter, X81699, (Helps et al. Eur. J. Biochem. 228:927-930, 1995). Furthermore, IPT-1 is 55% identical to human NaPi-3 over 625 amino acids (Magagnin et al. PNAS 90:5979-5983, 1993). The nucleotide sequence of Table 1 (SEQ ID NO:1) has about 78% identity (using FASTA) in 2242 nucleotide residues with bovine sodium dependent phosphate transporter, X81699, (Helps et al. Eur. J. Biochem. 228:927-930, 1995). Furthermore, IPT-1 is 58% identical to human NaPi-3 over 2270 nucleotides (Magagnin et al. PNAS 90:5979-5983, 1993). Thus, IPT-1 polypeptides and polynucleotides of the present invention are expected to have, inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1^a

5 CGGGCCAGGTTTCCAGGCTCGGCCGCCCTCCATCCCAGCACCTGCGGAGGGAGCGCTG
 ACCATGGCTCCCTGGCCTGAATTGGGAGATGCCAGCCCAACCCCGATAAGTACCTCGAA
 GGGGCCGAGGTGAGAGCCCACTGCCCTGATAAAAGCAAAGAGACCAACAAAACAGAT
 AACACTGAGGCACCTGTAACCAAGATTGAACCTCTGCCGTCTACTCCAGGCTACACTG
 10 ATAGATGAGCCCACTGAGGTGGATGACCCCTGGAACCTACCCACTCTTCAGGACTCGGGG
 ATCAAGTGGTCAGAGAGAGACACCAAGGGAAGATTCTCTGTTTCTTCCAAGGGATTGGG
 AGATTGATTTTACTTCTCGGATTCTCTACTTTTTCGTGTGCTCCCTGGATATTCTTAGT
 AGCGCCTTCCAGCTGGTTGGAGGAAAAATGGCAGGACAGTTCTTCAGCAACAGCTCTATT
 15 ATGTCCAACCCCTTTGTTGGGGCTGGTGATCGGGGTGCTGGTGACCGTCTTGGTGAGAGC
 TCCAGCACCTCAACGTCCATCGTTGTGAGCATGGTGTCTCTTCATTGCTCACTGTTCCG
 GCTGCCATCCCCATTATCATGGGGGCCAACATTGGAACGTCAATCACCAACACTATTGTT
 GCGCTCATGCAGGTGGGAGATCGGAGTGAGTTCAGAAGAGCTTTTGAGGAGCCACTGTC
 20 CATGACTTCTTCAACTGGCTGTCCGTGTTGGTGCTCTTGCCCGTGGAGGTGGCCACCCAT
 TACCTCGAGATCATAACCCAGCTTATAGTGGAGAGCTTCCACTTCAAGAATGGAGAAGAT
 GCCCCAGATCTTCTGAAAGTCATCACTAAGCCCTTCACAAAGCTCATTGTCCAGCTGGAT
 AAAAAAGTTATCAGCCAAATTGCAATGAACGATGAAAAAGCGAAAAACAAGAGTCTTGTC
 25 AAGATTTGGTGCAAACTTTTACCAACAAGACCCAGATTAACGTCAGTGTTCCTCGACT
 GCTAACTGCACCTCCCTTCCCTCTGTTGGACGGATGGCATCCAAAAGTGGACCATGAAG
 AATGTGACCTACAAGGAGAACATCGCCAAATGCCAGCATATCTTTCTGAATTTCCACCTC
 CCGGATCTTGCTGTGGGCACCATCTTGCTCATACTCTCCCTGCTGGTCTCTGTGGTTGC
 30 CTGATCATGATTGTCAAGATCCTGGGCTCTGTGCTCAAGGGGCAGGTGCGCACTGTCATC
 AAGAAGACCATCAACACTGATTTCCCTTTCCCTTTGCATGGTTGACTGGCTACCTGGCC
 ATCCTCGTCGGGGCAGGCATGACCTTCATCGTACAGAGCAGCTCTGTGTTCAAGTGGCC
 TTGACCCCTGATTGGAATCGGCGTGATAACCAATTGAGAGGGCTTATCCACTCAGGCTG
 35 GGCTCCAACATCCGCACCAACCACCGCCATCCTGGCCGCTTAGCCAGCCCTGGCAAT
 GCATTGAGGAGTTCACTCCAGATCGCCCTGTGCCACTTTTCTTCAACATCTCCGGCATC
 TTGCTGTGGTACCCGATCCCGTTCACTCGCTGCCATCCGCATGGCCAAGGGGCTGGCC
 AACATCTCTGCCAAGTATCGCTGGTTGCGCGTCTTCTACCTGATCATCTTCTTCTCTG
 40 ATCCCGCTGACGCTGTTTGGCCTCTCGCTGGCCGGCTGGCGGGTCTGGTTGGTGTGGG
 GTTCCCGTCTGTTTCATCATCATCCTGGTACTGTGCTCCGACTCCTSCAGTCTGCTGC
 CCACCGCTCTGCGGAAGAACTCCAGAACTGGAACCTTCTGCGCTGTGGTGGCGCTG
 45 CIGAAGCCCTGGGATGCCGCTGCTCTCAAGTTACCCGCTGCTTCCAGATCCGCTGCTGC
 TACTGCTGCCGCTGTGCTGCCGCGCTGCTGCTTGCTGTGTGGCTGCCCAAGTGTGCTGC
 CGCTGCAGCAAGTGTGCGAGGACTTGAGGAGGGGCGAGGAGGGGCAAGATGTCTGTGTC
 AAGGCTCCTGAGACCTTTGATAACATAACCATTAGCAGAGAGGCTCAGGGTGAGCTCCCT
 50 GCCTCGGACTCAAAGACCGAATGCACGGCCTGTAGGGGACGCCCCAGATTGTGAGGAT
 GGGGGGATGGTCTTTGAGTTTTCATGCTCTCTCCCTCCCACTTCTGCACCCCTTTCACC

55

ACCTCGAGGAGATTGCTCCCCATTAGCGAATGAAATTGATGCAGTCCTAAAAAAAAAAAA
 AAAAAAAA

^a A nucleotide sequence of a human IPT-1 (SEQ ID NO: 1).

Table 2^b

MAPWPELGDAQPNPDKYLEGAAGQQPTAPDKSKETNKTDNTEAPVTKEILLPSYSTATLI
 DEPTVEDDPWNLP TLQDSGIKWSEKDTKGKILCFFQIGIRLILLGLFLYFFVCSLDILSS
 AFQLVGGKMAQQFFSNSSIMSNPLGLVIGVLVTVLVQSSSTSTSI VVSMVSSSLT VRA
 AIP IIMGANIGTSITNTI VALMQVGRSEFRRAFAGATVHDFFNWLSVLVLLPVEVATHY
 LEIITQLIVESFHFKNGEDAPDLLKVITKPF TKLIVQLDKKVISQIAMNDEKAKNKS LVK
 FWCKTFTNKTIQINVTVPSTANCTSPSLCWT DGIQNWIMKNVTYKENTAKCQHI FVNFHLP
 DLAVGTILLIILSLVLGCLIMIVKTLGSLVKGVATVIKKTINTDFPFPAWLTGYLAI
 LVGAGMTFTVQSSSVFTSALTELGIGVITIERAYPLTLGSGNIGTTTTAILAALASPGNA
 LRSSLQIALCHFFFNISGILLWYPTFFTRLPTRMAGLGNISAKYRWFAVFYLI TFFFI :
 PLTVFGLSLAGWRVLVGVGVVVFII ILVLCRLRLQSRCPRVLPKKLQWNWFLPIWMRSL
 KPWDVAVSKFTGCFQMRCCYCCRVCCRCACLLCGCPKCCRCCKCEDLEEAQEGQDVPVK
 APETFDNITISREAQGEVPASDSKTECTAL

^b An amino acid sequence of a human IPT-1 (SEQ ID NO: 2).

One polynucleotide of the present invention encoding IPT-1 may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells of human small intestine and lung using the expressed sequence tag (EST) analysis (Adams, M.D., *et al. Science* (1991) 252:1651-1656 ; Adams, M.D. *et al., Nature*, (1992) 355:632-634; Adams, M.D., *et al., Nature* (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

The nucleotide sequence encoding IPT-1 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in Table 1 (nucleotide number 64 to 2136 of SEQ ID NO:1), or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

When the polynucleotides of the invention are used for the recombinant production of IPT-1 polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al., Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Further preferred embodiments are polynucleotides encoding IPT-1 variants comprising the amino acid sequence of IPT-1 polypeptide of Table 2 (SEQ ID NO:2) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination. Among the preferred polynucleotides of the present invention is contained in Table 3 (SEQ ID NO: 3) encoding the amino acid sequence of Table 4 (SEQ ID NO: 4).

Table 3^c

GAATTCGGCTTGCCACCAACGAGTTTCATCCTGAGCCTGCTGGTCTCTGTGTTTGCTGA
 TCATGATTGTCAAGATCCTGGGCTCTGTGCTAAGGGGCAGGTCCGCCACTGTTCATCAAGAA
 GACCATCAACACTGATTTCCCTTTCCCTTTGCAIGTTGACTGGCTACCTGGCCATCCT
 CGTCGGGGCAGGCATGACCTTCATCGTACAGAGCAGCTCTGTGTTACGTCGGCCTTGAC
 CCCCCTGATTGGAATCGGCGTGATAACCATTGAGAGGGCTTATCCACTCACGCTGGGCTC
 CAACATCGGCACCACCACCACCGCCATCCTGGCCGCTTAGCCAGCCCTGGCAATGCATT
 GAGGAGTTCACTCCAGATCGCCCTGTGCCACTTTTCTTCAACATCTCCGGCATCTTGCT
 GTGGTACCCGATCCCGTTCACTCGCCTGCCCATCCGCATGGCCAAGGGGCTGGGCAACAT
 CTCTGCCAAGTATCGCTGGTTCGCCGTCTTCTACCTGATCATCTTCTTCTCTGATCCC
 GCTGACGGTGTGTCCTCTGCTGGTTCCTCTCTACAAGCCGAATTCTGCAGATATCCA
 TCACACTGGCGGCCGCTCGAG

^c A partial nucleotide sequence of a human IPT-1 (SEQ ID NO: 3).

Table 4^d

KKTINTDFPFPAWLTGYXAIXVGAGMTFIVQSSSVETSALTPLIGIVITIERAYPLTL
 GSNIGTTTTAILAALASPGNALRSSQLIALCHFFFNISGILLWYPIPFTRLPIRMAKGLG
 NISAKYRWFVAFYLIIFFFLIPLTVFALCWLPLYK

^d A partial amino acid sequence of a human IPT-1 (SEQ ID NO: 4).

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO: 1 or a fragment thereof (including that of SEQ ID NO:3), may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding IPT-1 polypeptide and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to the IPT-1 gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

In one embodiment, to obtain a polynucleotide encoding IPT-1 polypeptide, including homologs and orthologs from species other than human, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO: 1 or a fragment thereof (including that of SEQ ID NO: 3), and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to those of skill in the art. Thus in another aspect, IPT-1 polynucleotides of the present invention further include a nucleotide sequence comprising a nucleotide sequence that hybridize under stringent condition to a nucleotide sequence having SEQ ID NO: 1 or a fragment thereof (including that of SEQ ID NO:3). Also included with IPT-1 polypeptides are polypeptide comprising amino acid sequence encoded by nucleotide sequence obtained by the above hybridization condition. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

Vectors, Host Cells, Expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY* (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL* (*supra*).

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If the IPT-1 polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If IPT-1 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered. IPT-1 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and/or purification.

Diagnostic Assays

This invention also relates to the use of IPT-1 polynucleotides for use as diagnostic reagents. Detection of a mutated form of IPT-1 gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of IPT-1. Individuals carrying mutations in the IPT-1 gene may be detected at the DNA level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled IPT-1 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science* (1985) 230:1242. Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. See Cotton et al., *Proc Natl Acad Sci USA* (1985) 85: 4397-4401. In another embodiment, an array of oligonucleotide probes comprising IPT-1 nucleotide sequence or fragments thereof

can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

5 The diagnostic assays offer a process for diagnosing or determining a susceptibility to chronic renal failure, end stage renal disease, uremic bone disease, and cancer, through detection of mutation in the IPT-1 gene by the methods described.

In addition, chronic renal failure, end stage renal disease, uremic bone disease, and cancer can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of
10 IPT-1 polypeptide or IPT-1 mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as an IPT-1 polypeptide, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease, particularly chronic renal failure, end stage renal disease, uremic bone disease, and cancer, which comprises:

- (a) a IPT-1 polynucleotide, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a IPT-1 polypeptide, preferably the polypeptide of SEQ ID NO: 2, or a fragment thereof; or
- (d) an antibody to a IPT-1 polypeptide, preferably to the polypeptide of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

Antibodies

The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the IPT-1 polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

Antibodies generated against the IPT-1 polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, MONOCLONAL ANTIBODIES AND CANCER THERAPY, pp. 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against IPT-1 polypeptides may also be employed to treat chronic renal failure, end stage renal disease, uremic bone disease, and cancer, among others.

Vaccin s

Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with IPT-1 polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immun response to protect said animal from chronic renal failure, end stage renal disease, uremic bone disease, and cancer, among others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering IPT-1 polypeptide via a vector directing expression of IPT-1 polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a IPT-1 polypeptide wherein the composition comprises a IPT-1 polypeptide or IPT-1 gene. The vaccine formulation may further comprise a suitable carrier. Since IPT-1 polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

The IPT-1 polypeptide of the present invention may be employed in a screening process for compounds which activate (agonists) or inhibit activation of (antagonists, or otherwise called inhibitors) the IPT-1 polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess identify agonist or antagonists from, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These agonists or antagonists may be natural or modified substrates, ligands, enzymes, receptors, etc., as the case may be, of the polypeptide of the present invention; or may be structural or functional mimetics of the polypeptide of the present invention. See Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991).

IPT-1 polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirous to find compounds and drugs which stimulate IPT-1 polypeptide on the one hand and which can inhibit the function of IPT-1 polypeptide on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions as chronic renal failure, end stage renal disease, uremic bone disease, and cancer. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions as chronic renal failure, end stage renal disease, uremic bone disease, and cancer.

In general, such screening procedures may involve using appropriate cells which express the IPT-1 polypeptide or respond to IPT-1 polypeptide of the present invention. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. Cells which express the IPT-1 polypeptide (or cell membrane containing the expressed polypeptide) or respond to IPT-1 polypeptide are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response. The ability of the cells which were contacted with the candidate compounds is compared with the same cells which were not contacted for IPT-1 activity.

Cell lines (transient or stable) expressing the recombinant protein are useful for establishing screening assays and for characterization of the protein. Phosphate uptake assays using these cells would be useful in the identification of inhibitors of phosphate transport. These assays could be either whole cell based or membrane based.

The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the IPT-1 polypeptide is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the IPT-1 polypeptide, using detection systems appropriate to the cells bearing the IPT-1 polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a IPT-1 polypeptide to form a mixture, measuring IPT-1 activity in the mixture, and comparing the IPT-1 activity of the mixture to a standard.

The IPT-1 cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of IPT-1 mRNA and protein in cells. For example, an ELISA may be

constructed for measuring secreted or cell associated levels of IPT-1 protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of IPT-1 (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The IPT-1 protein may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the IPT-1 is labeled with a radioactive isotope (eg 125I), chemically modified (eg biotinylated), or fused to a peptid sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. In addition to being used for purification and cloning of the receptor, these binding assays can be used to identify agonists and antagonists of IPT-1 which compete with the binding of IPT-1 to its receptors, if any. Standard methods for conducting screening assays are well understood in the art.

Examples of potential IPT-1 polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, enzymes, receptors, etc., as the case may be, of the IPT-1 polypeptide, e.g., a fragment of the ligands, substrates, enzymes, receptors, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for IPT-1 polypeptides; or compounds which decrease or enhance the production of IPT-1 polypeptides, which comprises:

- (a) a IPT-1 polypeptide, preferably that of SEQ ID NO:2;
- (b) a recombinant cell expressing a IPT-1 polypeptide, preferably that of SEQ ID NO:2;
- (c) a cell membrane expressing a IPT-1 polypeptide; preferably that of SEQ ID NO: 2; or
- (d) antibody to a IPT-1 polypeptide, preferably that of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

This invention provides methods of treating abnormal conditions such as, chronic renal failure end stage renal disease, uremic bone disease, and cancer, related to both an excess of and insufficient amounts of IPT-1 polypeptide activity.

If the activity of IPT-1 polypeptide is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as hereinabove described along with a pharmaceutically acceptable carrier in an amount effective to inhibit the function of the IPT-1 polypeptide, such as, for example, by blocking the binding of ligands, substrates, enzymes, receptors, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of IPT-1 polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous IPT-1 polypeptide may be administered. Typical embodiments of such competitors comprise fragments of the IPT-1 polypeptide.

In another approach, soluble forms of IPT-1 polypeptides still capable of binding the ligand in competition with endogenous IPT-1 polypeptide may be administered. Typical embodiments of such competitors comprise fragments of the IPT-1 polypeptide.

In still another approach, expression of the gene encoding endogenous IPT-1 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. See, for example, O'Connor, *JNeurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Alternatively, oligonucleotides which form triple helices with the gene can be supplied. See, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360. These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

For treating abnormal conditions related to an under-expression of IPT-1 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of compound which activates IPT-1 polypeptide, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of IPT-1 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For overview of gene therapy, see Chapter 20, *Gene Therapy*

and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and AP Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of IPT-1 polypeptides in combination with a suitable pharmaceutical carrier.

5 Formulation and Administration

Peptides, such as the soluble form of IPT-1 polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

This cDNA sequence can be expressed in heterologous systems (prokaryotic or eukaryotic) to allow high level protein expression and production. Recombinant protein produced in, and purified from, these systems is useful in screening assays to identify inhibitors of catalytic activity as well as a source of material for structural studies and antibody production. Cell lines (transient or stable) expressing the recombinant protein are useful for establishing screening assays and for characterization of the protein.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

Annex to the description

5

SEQUENCE LISTING

10

(1) GENERAL INFORMATION

15

(i) APPLICANT: SmithKline Beecham Corporation

(ii) TITLE OF THE INVENTION: A HUMAN SODIUM DEPENDENT PHOSPHATE
TRANSPORTER (IPT-1)

20

25

30

35

40

45

50

55

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: SmithKline Beecham, Corporate Intellectual
Property
- (B) STREET: Two New Horizons Court
- (C) CITY: Brentford
- (D) STATE: Middlesex
- (E) COUNTRY: United Kingdom
- (F) ZIP: TW8 9EP

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Diskette
- (B) COMPUTER: IBM Compatible
- (C) OPERATING SYSTEM: DOS
- (D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: TO BE ASSIGNED
- (B) FILING DATE: 23-SEP-1997
- (C) CLASSIFICATION: UNKNOWN

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 60/044,974
- (B) FILING DATE: 28-APR-1997

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: CONNELL, Anthony Christopher
- (B) REGISTRATION NUMBER: 5630
- (C) REFERENCE/DOCKET NUMBER: GH70006

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: +44 127 964 4395
- (B) TELEFAX: +44 181 975 6294
- (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2288 base pairs

(B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10	CGGGCCAGGT TTCCAGGCTC GGCCGCCGCC TCCATCCCAG CACCTGCGGA GGGAGCGCTG	60
	ACCATGGCTC CCTGGCCTGA ATTGGGAGAT GCCCAGCCCA ACCCCGATAA GTACCTCGAA	120
	GGGGCCGCAG GTCAGCAGCC CACTGCCCTT GATAAAAGCA AAGAGACCAA CAAAACAGAT	180
15	AACACTGAGG CACCTGTAAC CAAGATTGAA CTTCTGCCGT CCTACTCCAC GGCTACACTG	240
	ATAGATGAGC CCACTGAGGT GGATGACCCC TGGAACCTAC CCACTCTTCA GGACTCGGGG	300
	ATCAAGTGGT CAGAGAGAGA CACCAAAGGG AAGATTCTCT GTTTCTTCCA AGGGATTGGG	360
	AGATTGATTT TACTTCTCGG ATTTCTCTAC TTTTCTGTGT GCTCCCTGGA TATTCTTAGT	420
20	AGCGCCTTCC AGCTGGTTGG AGGAAAAATG GCAGGACAGT TCTTCAGCAA CAGCTCTATT	480
	ATGTCCAACC CTTTGTGTTGG GCTGGTGATC GGGGTGCTGG TGACCGTCTT GGTGCAGAGC	540
	TCCAGCACCT CAACGTCCAT CGTTGTCAGC ATGGTGTCTT CTTTATTGCT CACTGTTCCG	600
	GCTGCCATCC CCATTATCAT GGGGGCCAAC ATTGGAACGT CAATCACCAA CACTATTGTT	660
25	GCGCTCATGC AGGTGGGAGA TCGGAGTGAG TTCAGAAGAG CTTTTCAGAG AGCCACTGTC	720
	CATGACTTCT TCAACTGGCT GTCCGTGTTG GTGCTCTTGC CCGTGGAGGT GGCCACCCAT	780
	TACCTCGAGA TCATAACCCA GCTTATAGTG GAGAGCTTCC ACTTCAAGAA TGGAGAAGAT	840
	GCCCCAGATC TTCTGAAAGT CATCACTAAG CCCTTCACAA AGCTCATTGT CCAGCTGGAT	900
30	AAAAAAGTTA TCAGCCAAAT TGCAATGAAC GATGAAAAAG CGAAAAACAA GAGTCTTGTC	960
	AAGATTTGGT GCAAAACTTT TACCAACAAG ACCCAGATTA ACGTCACTGT TCCCTCGACT	1020
	GCTAACTGCA CCTCCCCTTC CCTCTGTTGG ACGGATGGCA TCCAAAAGTG GACCATGAAG	1080
	AATGTGACCT ACAAGGAGAA CATCGCCAAA TGCCAGCATA TCTTTGTGAA TTTCCACCTC	1140
35	CCGGATCTTG CTGTGGGCAC CATCTTGCTC ATACTCTCCC TGCTGGTCCT CTGTGGTTGC	1200
	CTGATCATGA TTGTCAAGAT CCTGGGCTCT GTGCTCAAGG GGCAGGTCGC CACTGTGCATC	1260
	AAGAAGACCA TCAACACTGA TTTCCCCTTT CCCTTTGCAT GGTTGACTGG CTACCTGGCC	1320
	ATCCTCGTCG GGGCAGGCAT GACCTTCATC GTACAGAGCA GCTCTGTGTT CACGTCGGCC	1380
40	TTGACCCCCC TGATTGGAAT CGGCGTGATA ACCATTGAGA GGGCTTATCC ACTCACGCTG	1440
	GGCTCCAACA TCGGCACCAC CACCACCGCC ATCCTGGCCG CCTTAGCCAG CCCTGGCAAT	1500
	GCATTGAGGA GTTCACTCCA GATCGCCCTG TGCCACTTTT TCTTCAACAT CTCGGGCATC	1560
	TTGCTGTGGT ACCCGATCCC GTTCACTCGC CTGCCATCC GCATGGCCAA GGGGCTGGGC	1620
45	AACATCTCTG CCAAGTATCG CTGGTTCGCC GTCTTCTACC TGATCATCTT CTTCTTCCTG	1680
	ATCCCCTGTA CGGTGTTTGG CCTCTCGCTG GCCGGCTGGC GGGTGCTGGT TGGTGTGCGG	1740
	GTTCCCGTCG TCTTCATCAT CATCCTGGTA CTGTGCCTCC GACTCCTGCA GTCTCGCTGC	1800
	CCACGCGTCC TGCCGAAGAA ACTCCAGAAC TGGAACCTCC TGCCGCTGTG GATGCGCTCG	1860
50	CTGAAGCCCT GGGATGCCGT CGTCTCCAAG TTCACCGCT GCTTCCAGAT GCGCTGCTGC	1920
	TACTGCTGCC GCGTGTGCTG CCGCGCGTGC TGCTTGCTGT GTGGCTGCCC CAAGTGCTGC	1980
	CGCTGCAGCA AGTGCTGCGA GGACTTGAG GAGGCGCAGG AGGGGCAGGA TGTCCCTGTC	2040
	AAGGCTCCTG AGACCTTGA TAACATAACC ATTAGCAGAG AGGCTCAGG TGAGGTCCCT	2100
55	GCCTCGGACT CAAAGACCGA ATGCACGGCC TTGTAGGGGA CGCCCCAGAT TGTCAGGGAT	2160

GGGGGGATGG TCCTTGAGTT TTGCATGCTC TCCTCCCTCC CACTTCTGCA CCCTTTCACC 2220
 ACCTCGAGGA GATTGCTCC CCATTAGCGA ATGAAATTGA TGCAGTCCTA AAAAAAAAAA 2280
 AAAAAAAAAA 2288

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 690 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Pro Trp Pro Glu Leu Gly Asp Ala Gln Pro Asn Pro Asp Lys
 1 5 10 15
 Tyr Leu Glu Gly Ala Ala Gly Gln Gln Pro Thr Ala Pro Asp Lys Ser
 20 25 30
 Lys Glu Thr Asn Lys Thr Asp Asn Thr Glu Ala Pro Val Thr Lys Ile
 35 40 45
 Glu Leu Leu Pro Ser Tyr Ser Thr Ala Thr Leu Ile Asp Glu Pro Thr
 50 55 60
 Glu Val Asp Asp Pro Trp Asn Leu Pro Thr Leu Gln Asp Ser Gly Ile
 65 70 75 80
 Lys Trp Ser Glu Arg Asp Thr Lys Gly Lys Ile Leu Cys Phe Phe Gln
 85 90 95
 Gly Ile Gly Arg Leu Ile Leu Leu Leu Gly Phe Leu Tyr Phe Phe Val
 100 105 110
 Cys Ser Leu Asp Ile Leu Ser Ser Ala Phe Gln Leu Val Gly Gly Lys
 115 120 125
 Met Ala Gly Gln Phe Phe Ser Asn Ser Ser Ile Met Ser Asn Pro Leu
 130 135 140
 Leu Gly Leu Val Ile Gly Val Leu Val Thr Val Leu Val Gln Ser Ser
 145 150 155 160
 Ser Thr Ser Thr Ser Ile Val Val Ser Met Val Ser Ser Ser Leu Leu
 165 170 175
 Thr Val Arg Ala Ala Ile Pro Ile Ile Met Gly Ala Asn Ile Gly Thr
 180 185 190
 Ser Ile Thr Asn Thr Ile Val Ala Leu Met Gln Val Gly Asp Arg Ser
 195 200 205
 Glu Phe Arg Arg Ala Phe Ala Gly Ala Thr Val His Asp Phe Phe Asn
 210 215 220

	Trp	Leu	Ser	Val	Leu	Val	Leu	Leu	Pro	Val	Glu	Val	Ala	Thr	His	Tyr	
	225					230					235					240	
5	Leu	Glu	Ile	Ile	Thr	Gln	Leu	Ile	Val	Glu	Ser	Phe	His	Phe	Lys	Asn	
				245					250						255		
	Gly	Glu	Asp	Ala	Pro	Asp	Leu	Leu	Lys	Val	Ile	Thr	Lys	Pro	Phe	Thr	
			260						265					270			
10	Lys	Leu	Ile	Val	Gln	Leu	Asp	Lys	Lys	Val	Ile	Ser	Gln	Ile	Ala	Met	
		275						280					285				
	Asn	Asp	Glu	Lys	Ala	Lys	Asn	Lys	Ser	Leu	Val	Lys	Ile	Trp	Cys	Lys	
	290						295					300					
15	Thr	Phe	Thr	Asn	Lys	Thr	Gln	Ile	Asn	Val	Thr	Val	Pro	Ser	Thr	Ala	
	305					310					315				320		
	Asn	Cys	Thr	Ser	Pro	Ser	Leu	Cys	Trp	Thr	Asp	Gly	Ile	Gln	Asn	Trp	
				325						330				335			
20	Thr	Met	Lys	Asn	Val	Thr	Tyr	Lys	Glu	Asn	Ile	Ala	Lys	Cys	Gln	His	
				340					345					350			
	Ile	Phe	Val	Asn	Phe	His	Leu	Pro	Asp	Leu	Ala	Val	Gly	Thr	Ile	Leu	
		355					360						365				
25	Leu	Ile	Leu	Ser	Leu	Leu	Val	Leu	Cys	Gly	Cys	Leu	Ile	Met	Ile	Val	
	370						375					380					
	Lys	Ile	Leu	Gly	Ser	Val	Leu	Lys	Gly	Gln	Val	Ala	Thr	Val	Ile	Lys	
	385					390					395				400		
30	Lys	Thr	Ile	Asn	Thr	Asp	Phe	Pro	Phe	Pro	Phe	Ala	Trp	Leu	Thr	Gly	
				405						410				415			
	Tyr	Leu	Ala	Ile	Leu	Val	Gly	Ala	Gly	Met	Thr	Phe	Ile	Val	Gln	Ser	
			420				425					430					
35	Ser	Ser	Val	Phe	Thr	Ser	Ala	Leu	Thr	Pro	Leu	Ile	Gly	Ile	Gly	Val	
			435				440					445					
	Ile	Thr	Ile	Glu	Arg	Ala	Tyr	Pro	Leu	Thr	Leu	Gly	Ser	Asn	Ile	Gly	
	450					455					460						
40	Thr	Thr	Thr	Thr	Ala	Ile	Leu	Ala	Ala	Leu	Ala	Ser	Pro	Gly	Asn	Ala	
	465					470					475				480		
	Leu	Arg	Ser	Ser	Leu	Gln	Ile	Ala	Leu	Cys	His	Phe	Phe	Phe	Asn	Ile	
				485						490				495			
45	Ser	Gly	Ile	Leu	Leu	Trp	Tyr	Pro	Ile	Pro	Phe	Thr	Arg	Leu	Pro	Ile	
				500					505				510				
	Arg	Met	Ala	Lys	Gly	Leu	Gly	Asn	Ile	Ser	Ala	Lys	Tyr	Arg	Trp	Phe	
			515				520					525					
50	Ala	Val	Phe	Tyr	Leu	Ile	Ile	Phe	Phe	Phe	Leu	Ile	Pro	Leu	Thr	Val	
	530					535					540						
	Phe	Gly	Leu	Ser	Leu	Ala	Gly	Trp	Arg	Val	Leu	Val	Gly	Val	Gly	Val	
	545					550				555				560			
55	Pro	Val	Val	Phe	Ile	Ile	Ile	Leu	Val	Leu	Cys	Leu	Arg	Leu	Leu	Gln	

565 570 575
 Ser Arg Cys Pro Arg Val Leu Pro Lys Lys Leu Gln Asn Trp Asn Phe
 580 585 590
 5 Leu Pro Leu Trp Met Arg Ser Leu Lys Pro Trp Asp Ala Val Val Ser
 595 600 605
 Lys Phe Thr Gly Cys Phe Gln Met Arg Cys Cys Tyr Cys Cys Arg Val
 610 615 620
 10 Cys Cys Arg Ala Cys Cys Leu Leu Cys Gly Cys Pro Lys Cys Cys Arg
 625 630 635 640
 Cys Ser Lys Cys Cys Glu Asp Leu Glu Glu Ala Gln Glu Gly Gln Asp
 645 650 655
 15 Val Pro Val Lys Ala Pro Glu Thr Phe Asp Asn Ile Thr Ile Ser Arg
 660 665 670
 Glu Ala Gln Gly Glu Val Pro Ala Ser Asp Ser Lys Thr Glu Cys Thr
 675 680 685
 20 Ala Leu
 690

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 621 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAATTCGGCT TGCCACCAAC GAGTTCATCC TGAGCCTGCT GGTCTCTGT GGTGCCTGA 60
 TCATGATTGT CAAGATCCTG GGCTCTGTGC TAAGGGGCAG GTCGCCACTG TCATCAAGAA 120
 40 GACCATCAAC ACTGATTTCC CCTTTCCCTT TGCATGGTTG ACTGGCTACC TGGCCATCCT 180
 CGTCGGGGCA GGCATGACCT TCATCGTACA GAGCAGCTCT GTGTTACGT CGGCCTTGAC 240
 CCCCCTGATT GGAATCGGCG TGATAACCAT TGAGAGGGCT TATCCACTCA CGCTGGGCTC 300
 CAACATCGGC ACCACCACCA CCGCCATCCT GGCCGCCTTA GCCAGCCCTG GCAATGCATT 360
 45 GAGGAGTTCA CTCCAGATCG CCCTGTGCCA CTTTTCCTT AACATCTCCG GCATCTTGCT 420
 GTGGTACCCG ATCCCGTTCA CTCGCCTGCC CATCCGCATG GCCAAGGGGC TGGGCAACAT 480
 CTCTGCCAAG TATCGCTGGT TCGCCGTCTT CTACCTGATC ATCTTCTTCT TCCTGATCCC 540
 GCTGACGGTG TTTGCCCTCT GCTGGTTGCC TCTCTACAAG CCGAATTCTG CAGATATCCA 600
 50 TCACACTGGC GGCCGCTCGA G 621

(2) INFORMATION FOR SEQ ID NO:4:

TCACACTGGC GGCCGCTCGA G

621

5

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 155 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

20 Lys Lys Thr Ile Asn Thr Asp Phe Pro Phe Pro Phe Ala Trp Leu Thr
 1 5 10 15
 Gly Tyr Xaa Ala Ile Xaa Val Gly Ala Gly Met Thr Phe Ile Val Gln
 20 25 30
 25 Ser Ser Ser Val Phe Thr Ser Ala Leu Thr Pro Leu Ile Gly Ile Gly
 35 40 45
 Val Ile Thr Ile Glu Arg Ala Tyr Pro Leu Thr Leu Gly Ser Asn Ile
 50 55 60
 30 Gly Thr Thr Thr Thr Ala Ile Leu Ala Ala Leu Ala Ser Pro Gly Asn
 65 70 75 80
 Ala Leu Arg Ser Ser Leu Gln Ile Ala Leu Cys His Phe Phe Phe Asn
 85 90 95
 35 Ile Ser Gly Ile Leu Leu Trp Tyr Pro Ile Pro Phe Thr Arg Leu Pro
 100 105 110
 Ile Arg Met Ala Lys Gly Leu Gly Asn Ile Ser Ala Lys Tyr Arg Trp
 115 120 125
 40 Phe Ala Val Phe Tyr Leu Ile Ile Phe Phe Phe Leu Ile Pro Leu Thr
 130 135 140
 Val Phe Ala Leu Cys Trp Leu Pro Leu Tyr Lys
 45 145 150 155

Claims

50

1. An isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the IPT-1 polypeptide of SEQ ID NO:2; or a nucleotide sequence complementary to said isolated polynucleotide.
- 55 2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:1 encoding the IPT-1 polypeptide of SEQ ID NO:2.
3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 80%

identical to that of SEQ ID NO:1 over its entire length.

4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: 1.

5. The polynucleotide of claim 1 which is DNA or RNA.

6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a IPT-1 polypeptide comprising an amino acid sequence, which has at least 80% identity with the polypeptide of SEQ ID NO:2 when said expression system is present in a compatible host cell.

7. A host cell comprising the expression system of claim 6.

8. A process for producing a IPT-1 polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.

9. A process for producing a cell which produces a IPT-1 polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a IPT-1 polypeptide.

10. A IPT-1 polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.

11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:2.

12. An antibody immunospecific for the IPT-1 polypeptide of claim 10.

13. A method for the treatment of a subject in need of enhanced activity or expression of IPT-1 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an agonist to said polypeptide; and/or
- (b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence that has at least 80% identity to a nucleotide sequence encoding the IPT-1 polypeptide of SEQ ID NO:2 over its entire length; or a nucleotide sequence complementary to said nucleotide sequence in a form so as to effect production of said polypeptide activity *in vivo*.

14. A method for the treatment of a subject having need to inhibit activity or expression of IPT-1 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an antagonist to said polypeptide; and/or
- (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said polypeptide; and/or
- (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said polypeptide for its ligand, substrate, or receptor.

15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of IPT-1 polypeptide of claim 10 in a subject comprising:

- (a) determining the presence or absence of mutation in the nucleotide sequence encoding said IPT-1 polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the IPT-1 polypeptide expression in a sample derived from said subject.

16. A method for identifying compounds which inhibit (antagonize) or agonize the IPT-1 polypeptide of claim 10 which comprises:

- (a) contacting a candidate compound with cells which express the IPT-1 polypeptide (or cell membrane expressing IPT-1 polypeptide) or respond to IPT-1 polypeptide; and
- (b) observing the binding, or stimulation or inhibition of a functional response; or comparing the ability of the

cells (or cell membrane) which were contacted with the candidate compounds with the same cells which were not contacted for IPT-1 polypeptide activity.

17. An agonist identified by the method of claim 16.

18. An antagonist identified by the method of claim 16.

19. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a IPT-1 polypeptide.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 98 30 2815 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	WO 96 34288 A (ELI LILLY AND CO.) 31 October 1996 * claims; p. 1-3, 13, 24, 28-30, 34-36, 39-45 *	1-12,15, 16,19	C12N15/12 C07K14/47
Y,D	HELPS, C. ET AL. : "Cloning, sequence analysis and expression of the cDNA encoding a sodium-dependent phosphate transporter ..." EUR. J. BIOCHEM., vol. 228, 1995, pages 927-930, XP002075653 * abstract; materials and methods *	1-12,15, 16,19	
Y	NI, B. ET AL.: "Molecular cloning, expression, and chromosomal localization of a human brain-specific ..." J. NEUROCHEM., vol. 66, 1996, pages 2227-2238, XP002075654 *abstract; materials and methods; fig. 2 *	1-12,15, 16,19	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07K C12N
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
MUNICH		27 August 1998	Hermann, R
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons S : member of the same patent family, corresponding document			

EPO FORM 1503 DS 02 (04/00)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 98 30 2815

Claim(s) searched completely:
1-12,16,19

Claim(s) searched incompletely:
15

Claim(s) not searched:
13,14

Reason for the limitation of the search (non-patentable invention(s)):

Article 52 (4) EPC - Diagnostic method practised on the human or animal body; Method for treatment of the human or animal body by therapy or surgery

Further limitation of the search

Claim(s) not searched:
17,18

Reason for the limitation of the search:

A compound is not sufficiently defined by a method for its identification. Technical information concerning the compound itself is necessary to allow a meaningful search.